



Extending the generality of leaf economic design principles in the cycads, an ancient lineage

Yong-Jiang Zhang^{1,2}*, Kun-Fang Cao³*, Lawren Sack⁴*, Nan Li⁵, Xue-Mei Wei⁵ and Guillermo Goldstein^{6,7}

¹Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan 666303, China; ²Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA; ³State Key Laboratory for Conservation and Utilization of Subtropical Agro-bioresources and College of Forestry, Guangxi University, Nanning, Guangxi 530004, China; ⁴Department of Ecology and Evolutionary Biology, University of California, 621 Charles E. Young Drive South, Los Angeles, CA 90095-1606, USA; ⁵National Cycad Germplasm Conservation Center, Fairylake Botanical Garden, Shenzhen and Chinese Academy of Sciences, 160 Xianhu Rd., Liantang, Shenzhen 518004, China; ⁶Department of Biology, University of Miami, PO Box 249118, Coral Gables, FL 33124, USA; ⁷Departamento de Ecologia Genetica y Evolucion, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Nuñez, Buenos Aires, Argentina

Authors for correspondence: Yong-Jiang Zhang Tel: +1 617 496 3580 Email: yongjiangzhang@oeb.harvard.edu

Kun-Fang Cao Tel: +86 771 3274251 Email: kunfangcao@gxu.edu.cn

Received: *13 May 2014* Accepted: *8 December 2014*

New Phytologist (2015) **doi**: 10.1111/nph.13274

Key words: Cycadales, *Cycas*, functional convergence, gymnosperms, leaf economic spectrum, leaf hydraulic conductance, photosynthetic capacity, trade-off.

Introduction

Summary

• Cycads are the most ancient lineage of living seed plants, but the design of their leaves has received little study. We tested whether cycad leaves are governed by the same fundamental design principles previously established for ferns, conifers and angiosperms, and characterized the uniqueness of this relict lineage in foliar trait relationships.

• Leaf structure, photosynthesis, hydraulics and nutrient composition were studied in 33 cycad species from nine genera and three families growing in two botanical gardens.

• Cycads varied greatly in leaf structure and physiology. Similarly to other lineages, light-saturated photosynthetic rate per mass (A_m) was related negatively to leaf mass per area and positively to foliar concentrations of chlorophyll, nitrogen (N), phosphorus and iron, but unlike angiosperms, leaf photosynthetic rate was not associated with leaf hydraulic conductance. Cycads had lower photosynthetic N use efficiency and higher photosynthetic performance relative to hydraulic capacity compared with other lineages.

• These findings extend the relationships shown for foliar traits in angiosperms to the cycads. This functional convergence supports the modern synthetic understanding of leaf design, with common constraints operating across lineages, even as they highlight exceptional aspects of the biology of this key relict lineage.

Cycads are the most ancient living seed plant lineage (Brenner et al., 2003), considered as living fossils because of the morphological similarity between fossil and extant species. The oldest known cycad fossil dates to the Paleozoic (Mamay, 1969; Gao & Thomas, 1989), and the group reached dominance in the Mesozoic, the age of dinosaurs. Cycads show high diversity in morphology and structure, with leaves that are typically compound and possessing a thick cuticle (Norstog & Nicholls, 1997; Whitelock, 2002). Cycads vary greatly in their leaf vein system; some have single midveins in leaflets, others have midveins with dichotomizing lateral veins, and yet others have multiple parallel veins as in ferns (Stevenson et al., 1996). Leaf veins usually contain tracheids, but primitive vessels have also been found in some Cycas species (Huang & Zhang, 1999; Sack & Scoffoni, 2013). The leaf structural and physiological trait variation, and its association with light-saturated photosynthetic rate have not been quantified for cycads although these have been well established for angiosperms of a wide range of ecosystems and taxonomic groups. Indeed, it is not well-known whether cycad leaves show strong variation in physiology, or whether that variation would be constrained by similar principles as in angiosperms. In contrast to typical angiosperms, cycads have no response in stomatal density to elevated CO_2 (Haworth *et al.*, 2011) suggesting that they may have a relatively inflexible leaf design and physiology compared with angiosperms. The aim of this study was to assess for cycads the association of traits with photosynthetic rate, as expected based on studies of angiosperms, conifers, ferns and mosses (Fig. 1), including nutrient composition, leaf structure, stomatal conductance and leaf hydraulics.

The first trait tested for association with leaf photosynthetic rate was nutrient concentration. Nitrogen (N) is a crucial component of photosynthetic proteins, especially Rubisco (Field & Mooney, 1986). Total Rubisco activity and amount increase as leaf N increases (Sage *et al.*, 1987), and across angiosperm, conifer and fern species, leaf photosynthetic rate tends to correlate with leaf N on an area and/or mass basis (Fig. 1) (Field &

^{*}These authors contributed equally to this work.



Fig. 1 Relationships among leaf nitrogen (N) concentration (N_m), leaf mass per area (LMA, or canopy mass per area for mosses) and light-saturated photosynthetic rate per mass (A_m). Closed black circles, Glopnet (mainly angiosperms but also with conifers and three cycads) (Wright *et al.*, 2004); closed green circles, mosses (Waite & Sack, 2010); open circles, ferns (Karst & Lechowicz, 2007).

Mooney, 1986; Reich *et al.*, 1997; Wright *et al.*, 2004; Karst & Lechowicz, 2007). Other nutrients such as phosphorus (P), potassium (K), iron (Fe), sulfur (S) and zinc (Zn), are also important components of the photosynthetic apparatus and reactions (e.g. Randall & Bouma, 1973; Spiller & Terry, 1980; Ares *et al.*, 2009). However, perhaps due to a bias based on the much greater abundance of N and P, to our knowledge no interspecific relationships between leaf photosynthetic rate and nutrients other than N and P have been tested in any lineage.

In angiosperms, the effect of leaf structure on photosynthetic rate is well recognized (Reich et al., 1998; Niinemets & Sack, 2006). Across angiosperm and conifer species, light-saturated photosynthetic rate per leaf dry mass (A_m) and leaf N concentration per mass (Nm) correlate negatively with leaf mass per area (LMA) across a wide range of ecosystems (Fig. 1), typically independently of climate (Reich et al., 1997; Wright et al., 2004). These relationships, known as 'leaf economic spectrum' (LES) relationships, arise when photosynthetic rate per leaf area (A_a) and leaf N per area (Na) are partially or fully independent of LMA. The conversion of A_a to A_m by dividing by LMA thus results in LES relationships by mathematical necessity (Lloyd et al., 2013; Osnas et al., 2013). Yet, these relationships do reflect a physical influence of leaf structure on photosynthetic rate per mass (Sack et al., 2013; Westoby et al., 2013). Plants with higher LMA, due to thicker and/or denser leaves, tend to have a longer diffusion path and greater diffusion resistance from stomata to chloroplasts. Additionally, greater mass 'dilution' of cytoplasmic and chloroplast components by nonmetabolic components such as cell walls and nonstructural carbohydrates (Parkhurst, 1994; Reich et al., 1997, 1998; Roderick et al., 1999a,b; Shipley et al., 2006) would lead to a lower $A_{\rm m}$. The negative correlation between LMA and N_m is also due to the same mass dilution effect (Wright et al., 2004; Sack et al., 2013; Westoby et al., 2013). We expected that this mass dilution effect also exists for other nutrients and mediates the photosynthetic nutrient use

efficiency. Recently, a hypothesis that LMA and the LES relationships would be structurally determined by leaf vein density has been proposed (Blonder *et al.*, 2011, 2013, 2014). However, a recent test found that LMA was uncorrelated with leaf vein density across 275 phylogenetically diverse angiosperm species, and within eight of the nine individual families for which sufficient data were available (Sack *et al.*, 2013, 2014). Like conifers, which show LES relationships while having only a single central vein in their needles (Reich *et al.*, 1998), cycads provide a key system for determining whether LES relationships can arise in leaves that lack minor veins within their lamina.

The third trait tested for association with photosynthetic rate was xylem water transport efficiency. Leaf or stem hydraulic conductance has been found to be closely associated with leaf transpiration rate and maximum photosynthetic rate within the same species under different environmental conditions, within several sets of angiosperm species, and across representatives of many lineages (Kuppers, 1984; Meinzer & Grantz, 1990; Sober, 1997; Mencuccini, 2003; Santiago et al., 2004; Brodribb et al., 2007; Campanello et al., 2008; Zhang & Cao, 2009). These data, and manipulative experiments, have indicated that hydraulic conductance places a constraint on stomatal conductance (g_s) and on photosynthetic gas exchange. The leaf hydraulic conductance (K_{leaf}) plays a particularly important role because the leaf is a bottleneck in the whole-plant water transport pathway (Sack & Frole, 2006; Brodribb et al., 2007). Although K_{leaf} is well-recognized as a key determinant of A_a across species in very diverse species sets, the relationships of gas exchange with K_{leaf} may be climate- or habitat-specific, as they would tend to shift for optimality between habitats (Sack et al., 2005; Feild et al., 2011b; Sack & Scoffoni, 2013). Additionally, a negative relationship between mass-normalized K_{leaf} ($K_{\text{leaf-m}}$) and LMA has been reported for angiosperms (Simonin et al., 2012). Given the negative relationship of $A_{\rm m}$ to LMA, these relationships may result in a positive association of $A_{\rm m}$ and $K_{\rm leaf-m}$.

All of these previous findings represent a modern synthetic understanding that leaf structure, nutrient composition, water flux and photosynthetic rates are all related (e.g. Wright et al., 2003; Cramer et al., 2009), and are subject to the general principles of leaf design. However, to confirm the generality of these principles of leaf design, trait relationships need to be evaluated across different ecosystems and taxonomic groups, including lineages other than angiosperms. Cycads represent an ideal model for such a test, given the diversity of extant species, and the earlier origin and previous global dominance of this lineage. We studied cycads in two botanical gardens with similar climates to determine trait differences arising from genetic rather than plastic expression across natural habitats (Monson, 1996). We tested the hypotheses that photosynthetic performance is related to hydraulics, nutrient composition and leaf structure in cycads, and the relationships among functional traits should be similar in cycads to those found for other seed plants. We aimed to extend the understanding and generality of fundamental leaf trait relationships by focusing on the modern representatives of this ancient lineage and to characterize the uniqueness of cycads in foliar trait relationships.

Materials and Methods

Study site and plant material

This research was carried out in the Cycad Garden at Xishuangbanna Tropical Botanical Garden (XTBG; 21°41'N, 101°25'E, elevation 570 m) in southern Yunnan Province and the National Cycad Germplasm Conservation Center at Fairylake Botanical Garden (FBG; 22°34'N, 114°10'E, elevation 100-130 m) in Shenzhen, Guangdong Province, southern China. Mean annual temperature at XTBG is 21.7°C, and mean annual precipitation is 1560 mm with 80-85% occurring in the May-October rainy season (40-yr average, 1959–1998, data from Xishuangbanna Station for Tropical Rain Forest Ecosystem Studies). The soil of the Cycad Garden at XTBG is sandy alluvium with pH of 5.0-6.0; its total nutrient composition is $1.12 \text{ mg g}^{-1} \text{ N}$, 0.51 mg g^{-1} P, 8.49 mg g⁻¹ K, 8.33 mg g⁻¹ calcium (Ca), 2.89 mg g⁻¹ magnesium (Mg), 23.04 mg g⁻¹ Fe, 0.56 mg g^{-1} manganese (Mn), 73.25 mg kg⁻¹ Zn, and 22.0 mg kg⁻¹ Cu (0–20-cm depth). Soil hydrolyzable N, available P and available K are 66.5, 13.4 and 85.8 mg kg^{-1} , respectively. Mean annual temperature at FBG is 22.4°C, and mean annual precipitation is 1933 mm falling mostly in the May-October rainy season (30-yr average, 1971-2000, data from Shenzhen Whether Station). The soil of the National Cycad Germplasm Conservation Center at FBG is yellow soil with pH of 5.5–7.0, containing $1.25 \text{ mg g}^{-1} \text{ N}$, $0.52 \text{ mg g}^{-1} \text{ P}, \ 16.1 \text{ mg g}^{-1} \text{ K}, \ 1.20 \text{ mg g}^{-1} \text{ Ca}, \ 2.82 \text{ mg g}^{-1}$ Mg, 30.13 mg g^{-1} Fe, 0.42 mg g^{-1} Mn, 51.4 mg kg^{-1} Zn, and 9.43 mg kg^{-1} Cu (0–20-cm depth) in total. Soil hydrolyzable N, available P and available K are 115, 132 and 141 mg kg⁻¹, respectively. The physiological measurements at XTBG were performed in the rainy season 2009, and the measurements at FBG were performed in the rainy season 2011.

Thirty-three cycad species with at least three healthy and mature individuals suitable for physiological measurements were selected for the present study, with 10 from genus Cycas (family Cycadaceae), 19 from genera Ceratozamia, Dioon, Encephalartos, Lepidozamia, Macrozamia and Zamia (family Zamiaceae), and three from genera Bowenia and Stangeria (family Stangeriaceae) (Supporting Information Table S1). These species vary highly in native habitats (Table S1). Based on their native habitats, the species were classified into three classes: native to dry habitats, moist habitats, or both dry and moist habitats (for habitat information see Table S1). The genus Cycas is a sister lineage to all other cycads based on the phylogeny of cycads inferred from morphological characters or gene sequences (Brenner et al., 2003; Chaw et al., 2005; Zgurski et al., 2008). A fossil-calibrated phylogeny dates the Cycas lineage to the Mesozoic, whereas the lineage leading to Zamia dates to the Cenozoic (Nagalingum et al., 2011). Zamia has evenly spaced, longitudinally parallel, dichotomizing veins in leaflets whereas Cycas only has a single middle vein (Stevenson *et al.*, 1996). Leaf hydraulic conductance (K_{leaf}) was measured for 23 species, given difficulties in removing the effects of mucilage on K_{leaf} measurement for the remaining 10 species. Sun-exposed individuals were used for physiological measurements except for three shade-tolerant species which were

cultivated under a tree overstorey (*B. spectabilis, B. serrulata*) or in a shade house (*S. eriopus*). Four species were studied in both XTBG and FBG (*Cycas szechuanensis, C. panzhihuaensis, Z. furfuracea, E. gratus*), and these four species were entered twice in the dataset. Six mature leaves from three to six individuals per species were selected for the measurements. Leaflets were chosen as the fundamental unit for measurement and the middle section of the leaves were used for physiological measurements such as gas exchange and nutrient compositions. Leaflets were used, as these are most analogous to angiosperm simple leaves or leaflets, typically measured in studies of functional traits.

LMA, leaf thickness and density

Leaf thickness was averaged from values taken with vernier calipers at five places on each fresh leaflet. Leaf area was determined (LI-3000A area meter; Li-Cor, Lincoln, NE, USA), then the leaves were oven-dried at 70°C to constant mass, and weighed, and LMA was determined as leaf dry mass divided by leaf area (g m⁻²). Leaf density (mass/volume) was determined as LMA divided by leaf thickness (Witkowski & Lamont, 1991).

Gas exchange measurements

The light-saturated net CO₂ assimilation rate per area (A_a ; μ mol m⁻² s⁻¹) and stomatal conductance (g_s ; mol m⁻² s⁻¹) were measured using a portable photosynthetic system (LI-6400; Li-Cor). Leaves were measured on sunny days between 08:30 and 10:30 h solar time, at a photosynthetic photon flux density of 1500 μ mol m⁻² s⁻¹, and ambient temperature (28–35°C) and CO₂ concentration. Relative humidity ranged from 40% to 63%. Leaf temperatures ranged from 28 to 36°C at XTBG, and 27 to 32°C at FBG. Light-saturated net CO₂ assimilation rate per mass (A_m ; nmol g⁻¹ s⁻¹) was determined as A_a divided by LMA. A_a and g_s relative to hydraulic supply, representing leaf photosynthetic performance relative to investment in leaf hydraulic system (A_K and g_K ; mmol MPa mol⁻¹ and mol MPa mol⁻¹, respectively), were calculated as A_a and g_s , respectively, divided by K_{leaf-a} (mmol m⁻² s⁻¹ MPa⁻¹; see later).

Nutrient and chlorophyll concentrations

Leaves with rachis removed were oven-dried at 70°C for 48 h and analyzed for C, N, P, K, Ca, Zn, Fe and S. Leaf C and N concentrations were determined using a Vario MAX CN auto element analyzer (Elementar Analysensysteme, Hanau, Germany). After the samples were digested with concentrated HNO_3 – $HClO_4$, leaf P, K, Ca, Zn, Fe and S concentrations were measured using an inductively coupled plasma atomic-emission spectrometer (IRIS Advantage-ER; Thermo Jarrell Ash Corporation, Franklin, MA, USA). Nutrient concentrations per unit area were calculated by multiplying mass-based nutrient concentrations by LMA. Photosynthetic N use efficiency (A_N) and P use efficiency (A_P) were determined as A_m divided by mass-based concentrations of N and P (N_m and P_m) respectively. Leaf chlorophyll concentration was measured with a chlorophyll meter (SPAD 502; Minolta Corp., Osaka, Japan) for six sun-exposed mature leaves from three to six individuals per species. For each leaf, 10 measurements across the lamina were averaged (Chl*a*; in SPAD units, using a SPAD meter; SPAD-502; Minolta Corp.). Previous studies reported strong relationships between SPAD measurements and total extracted Chl*a* + *b* per area when leaves varied in nutrient composition, thickness and surface features (Marquard & Tipton, 1987; Singha & Townsend, 1989; Fanizza *et al.*, 1991; Manetas *et al.*, 1998). Furthermore, when multiple species have been compared, the majority tended to have similar relationships (Marquard & Tipton, 1987; Gratani, 1992; Markwell *et al.*, 1995; Castelli *et al.*, 1996; Uddling *et al.*, 2007; Marenco *et al.*, 2009), suggesting that SPAD measurements can be considered as a proxy for extractable chlorophyll. Chlorophyll concentration per mass (Chl_m) was determined as Chl*a* divided by LMA.

Leaf hydraulic conductance

Leaf hydraulic conductance (K_{leaf}) was measured using the evaporative flux method (Sack et al., 2002; Brodribb et al., 2007; Sack & Scoffoni, 2012). Leaves were cut from the plants in the late afternoon, recut immediately under water and placed with cut ends underwater and covered with black plastic bags to rehydrate overnight. Measurements of K_{leaf} were made on central segments of leaves, excised underwater, including rachis and 4-18 leaflets rather than on individual leaflets because petiolules of the leaflets were very short or nonexistent. The leaf segment was held horizontal in a frame strung with fishing line, above a fan, and under full sunlight to drive transpiration, with the cut end of the rachis connected to water-filled tubing downstream of a water pressuredrop flow meter with two calibrated pressure transducers (Melcher et al., 2012) logging to a computer, and the transpiration flow rate (E) was monitored. When a steady-state E was reached (i.e. a stable value maintained for at least 3 min), the leaf segment was removed into a ziplock bag that had been exhaled into, to prevent dehydration. Leaf water potential (Ψ_L) was determined after at least 30 min equilibration, using a pressure chamber (SKPM 1400; Skye Instruments Ltd, Lladrindod Wells, UK), and leaf area was measured (LI-3000A area meter; Li-Cor). The leaf area-normalized leaf hydraulic conductance $(K_{\text{leaf-a}}; \text{ mmol m}^{-2} \text{ s}^{-1} \text{ MPa}^{-1})$ was calculated as the steady-state $E'-\Psi_{\rm I}$, normalized by leaf area. Leaf temperature during measurement was monitored with a T-type thermal couple logging to a data logger (CR1000; Campbell Scientific, Logan, UT, USA), and $K_{\text{leaf-a}}$ was then corrected for the effects of temperature on the viscosity of water by standardizing to 20°C (Sack et al., 2002; Brodribb et al., 2007). The $K_{\text{leaf-a}}$ reported is the maximum, determined by plotting $K_{\text{leaf-a}}$ against Ψ_{L} and extrapolating to Ψ_{L} of zero (Brodribb et al., 2007). Maximum mass-normalized leaf hydraulic conductance (Kleaf-m, leaf water supply per dry mass; mol s⁻¹ g⁻¹ MPa⁻¹) was determined as $K_{\text{leaf-a}}$ divided by LMA.

Data analyses

Statistical analyses were applied using SPSS V21 (IBM Corp., Armonk, NY, USA). To examine trait-trait linkages, we tested relationships hypothesized a priori (see the Introduction section and Fig. 1). We analyzed the relationship between functional traits using linear or nonlinear regression according to which best approximated the structure of the relationship. Equal variances of the variables were tested and one-way ANOVAs were used to test trait differences among cycad genera. The differences in slope and intercept of bivariate relationships between cycads and plants from the global datasets (Wright *et al.*, 2004) or an angiosperm K_{leaf} dataset (Simonin et al., 2012) were tested with standardized major axis tests using SMATR v2.0 (Warton et al., 2006). The study site effect on leaf gas exchange characteristics (A_a , g_s and A_m) was tested by a two-way ANOVA with garden as the fixed factor and species as the random factor using four species studied in both XTBG and FBG (C. szechuanensis, C. panzhihuaensis, Zamia furfuracea, Encephalartos gratus). A principal components analysis (PCA), (Pearson, 1901) was performed to summarize the joint variation of the functional traits (see Table S2) for the 33 species, with mean trait values of each species used for the analyses.

Results

Variation of leaf traits among diverse cycad species

The cycad species varied strongly in leaf gross morphology and shape (Fig. 2), as well as in photosynthetic rate, and stomatal and hydraulic conductance (Tables 1, S3). Stomatal conductance (g_s) varied 10-fold (from 0.03 to 0.32 mol m⁻² s⁻¹), whereas leaf hydraulic conductance per mass ($K_{\text{leaf-m}}$) varied 13-fold (from 12.1 to 157 mol s⁻¹ g⁻¹ MPa⁻¹; Table S3).

Cycad species also varied strongly in nutrient concentrations and LMA, leaf thickness and leaf density (Tables 1, S3). N_a and P_a varied four- to five-fold, and the area based concentrations of other nutrients varied four-fold (S_a) to 20-fold (Ca_a). N_m and P_m varied across species three- to four-fold, whereas mass based concentrations of other nutrients varied 1.1-fold (C_m) to 20-fold (Fe_m). LMA, leaf thickness and leaf density varied five- to sevenfold across the species (Tables 1, S3). Cycad species also varied strongly in photosynthetic use-efficiency in relation to nitrogen, stomatal conductance and hydraulic conductance (Table 1).

The two genera of shade-tolerant species in family Stangeriaceae (*Bowenia* and *Stangeria*) showed significantly lower g_s , A_a and LMA than other cycad genera, whereas other genera did not differ significantly on average in any measured trait (P > 0.05, one-way ANOVA). Further, no significant difference was found between family Cycadaceae and family Zamiaceae in any measured trait (one-way ANOVA). No study site effects (difference between two gardens) on A_a , g_s and A_m were detected (P > 0.05, two-way ANOVA).

Coordination of leaf traits among diverse cycad species

The leaf economic trait relationships that were hypothesized based on those established for angiosperms were also very strong across the cycad species, and there were stronger trait relationships for mass-based than area-based light-saturated CO₂ assimilation rate. The A_a was positively related to g_s ($r^2 = 0.74$,





Fig. 2 Diversity in leaf form for 15 of the 32 cycad species studied.

P < 0.001, relationship not shown). Additionally, $A_{\rm m}$ was negatively related to LMA, and positively related to Chl_m, N_m, P_m, Fe_m (Fig. 3), K_m and S_m (relationships not shown). The relationship between $A_{\rm m}$ and Fe_m was found when data from each site were analyzed separately (P < 0.001). These relationships were significant even without including the two Zamia species (Z. fischeri, Z. vazquezii) that had substantially higher $A_{\rm m}$ (and in the case of Z. fischeri, substantially higher Fe_m) than other species (Table S3; dashed lines in Fig. 3).

In general, many cycad leaf traits were related to LMA, and particularly mass-based traits (Fig. 4). Photosynthetic N and P

use efficiencies (A_N and A_P), N_m , Chl_m , P_m , S_m and Fe_m were negatively related to LMA (Fig. 4). Although no statistical association was found between K_{leaf-a} and A_a (Fig. 5a), a significant relationship was found when they were expressed on a mass basis, that is, between K_{leaf-m} and A_m (Fig. 5b).

Comparison of trait ranges with those in previous studies of cycads and other lineages

Our data indicated a very strong physiological diversity among the cycads and similar trait relationships as previously reported

Table 1 Comparison of cycads in this study with Glopnet (Wright *et al.*, 2004), data spanning plant lineages, mosses and ferns in the mean and range of light-saturated photosynthetic rate per area (A_a), light-saturated photosynthetic rate per mass (A_m), photosynthetic nitrogen (N) use efficiency (A_N), light-saturated photosynthetic rate over leaf hydraulic conductance (A_k), N concentration per area (N_a), N concentration per mass (N_m), photosynthetic (N_m), p

	A _a	A _m	A _N	A _K	N _a	N _m	P _a	P _m	$K_{\text{leaf-a}}$	LMA
Cycads										
Mean	9.34	59.5	2.95	2.07	3.38	2.00	0.20	0.12	5.44	178
Minimum	3.15	23.6	1.42	0.61	1.52	1.22	0.08	0.07	2.50	50
Maximum	14.1	179	6.06	3.98	6.57	3.01	0.39	0.24	10.1	323
Glopnet										
Mean	11.5	128	6.31		1.94	1.94	0.13	0.11		128
Minimum	1.00	4.80	0.63		0.26	0.25	0.02	0.01		14
Maximum	42.0	662	25.5		9.14	6.36	0.88	0.60		1510
Sampling across p	olant lineages									
Mean	8.16			1.12					8.11	
Minimum	0.89			0.35					0.50	
Maximum	19.0			3.03					21.1	
Mosses										
Mean	0.93	8.11	0.01		0.08	0.80				13
Minimum	0.59	3.31	0.01		0.03	0.24				6
Maximum	1.41	14.0	0.03		0.12	1.51				31
Ferns										
Mean	4.50	67.5	2.88	1.30	1.64	1.75	0.08	0.09	2.93	90
Minimum	1.10	9.40	0.86	0.67	0.53	0.80	0.03	0.04	0.50	37
Maximum	12.3	115	5.29	2.20	3.54	3.29	0.20	0.13	5.23	158

Data source: cycads, 33 species (present study); Glopnet, 2548 species mainly angiosperms but also including conifers and including three cycads (Wright *et al.*, 2004); data spanning plant lineages, 12 angiosperms, 17 gymnosperms, six ferns, four bryophytes, five lycophytes (Brodribb *et al.*, 2007); Mosses, 10 species (Waite & Sack, 2010); ferns, 19 species (Wright *et al.*, 2004; Brodribb *et al.*, 2007).

for other lineages. The wide range of values for $K_{\text{leaf-a}}$ and A_a for the 33 cycad species studied encompassed the full range of values previously reported for conifers (Fig. 5a). However, the A_a of cycads were at the high end of the A_a range of conifers and had a higher median (Fig. 5c), whereas $K_{\text{leaf-a}}$ of cycads showed similar range and median relative to conifers (Fig. 5d). This helps explain the high A_K values of cycads (Table 1). The A_a of two cycad species (*C. cairnsiana and C. media*) previously measured by Brodribb *et al.* (2007) were close to the mean values of the cycads in the present study, whereas $K_{\text{leaf-a}}$ of those two species were close to the highest value found in this study.

By contrast, in several features, cycads showed lower maximum physiological capacities than angiosperms. Thus, the maximum $A_{\rm a}$, $A_{\rm m}$ and $A_{\rm N}$ in cycads were much lower than those in a global database consisting mainly of angiosperm species (Glopnet; Table 1). The mean A_a of cycads was very close to the Glopnet mean and slightly higher than the mean for a dataset of representatives from several plant lineages (angiosperms, conifers, ferns, bryophytes and lycophytes), and much higher than values compiled for mosses and ferns (Table 1). The mean LMA, and massand area-based concentrations of N and P for cycads were higher than the Glopnet mean, and also higher than those compiled for mosses or ferns. Consequently, the mean $A_{\rm m}$ and $A_{\rm N}$ values of cycads were much lower than the Glopnet mean (Wright et al., 2004), slightly higher than that of ferns, and much higher than that of mosses. The mean $K_{\text{leaf-a}}$ of cycads was low relative to other plant lineages, but, the $A_{\rm K}$ of cycads was higher than the mean for a dataset with representatives spanning all lineages, and higher than that compiled for ferns.

The bivariate relationships among LMA, A_m and N_m in cycads were generally consistent with the relationships for the species in Glopnet. No statistical differences in slope were found between the two datasets, although the relationships differed significantly in their intercepts (Fig. 6). Cycads tended to have a higher N_m at a given LMA, and a lower A_m at a given N_m , indicating lower photosynthetic N use efficiency compared with the global mean (Fig. 6b,c).

The relationship between K_{leaf} and LMA in cycads generally followed the relationship for species in a combined dataset mainly for angiosperms (Simonin *et al.*, 2012). No relationship was found between $K_{\text{leaf-a}}$ and LMA (Fig. 7a), whereas $K_{\text{leaf-m}}$ was significantly negatively related to LMA (Fig. 7b). No difference in slope was found between the two datasets, although the intercepts differed significantly (P < 0.01; Fig. 7b). The intercepts for log($K_{\text{leaf-m}}$) vs log(LMA) of cycads and angiosperms were 5.31 and 4.84 mol s⁻¹ g⁻¹ MPa⁻¹, respectively.

In a PCA analysis, the first axis, which explained 32% of the total variation, represented the variation in investment in leaf mass structure, and thus with leaf composition and physiology. The first PCA axis was positively related to LMA, leaf thickness, C_m and Chl*a*, and negatively correlated with A_m , Chl_m, A_N , A_P , $K_{\text{leaf-m}}$, N_m, S_m, Fe_m, P_m, Zn_m, K_m and $K_{\text{leaf-a}}$ (Fig. 8a). The second axis, which explained 21% of the total variation, represented the variation in stomatal opening for gas exchange and water use efficiency; this axis was positively correlated with g_s , A_a , $K_{\text{leaf-a}}$, g_K and A_K , and negatively correlated with water use efficiency (A/g). Zamia fischeri and Z. vazquezii were separated far from other species along the first axis, whereas shade-tolerant species were



Fig. 3 Light-saturated photosynthetic rate per mass (A_m) in relation to leaf mass per area (LMA), chlorophyll concentration per mass (Chl_m) and leaf nutrient concentrations per mass (N_m , P_m , Fe_m) across cycads in Xishuangbanna Tropical Botanical Garden (XTBG, green circles) and Fairylake Botanical Garden (FBG, red circles). Points are means \pm SE. Exponential (a, e) or linear (b–d) regressions were fitted to the data. The regressions do not include shade-tolerant species (open circles). The linear regressions indicated by the broken lines do not include *Zamia fischeri* and *Z. vazquezii*.

separated from other species along the second axis (Fig. 8b). The first and second axes could not separate cycads from FBG and XTBG, but most plants in XTBG fell into the upper two quadrants (Fig. 8b). Species native in dry habitats were not separated from those native in moist habitats or those found in both dry and moist habitats (Fig. 8c).

Discussion

Cycads have previously been described as highly diverse in their overall morphology (Whitelock, 2002). One major novel finding of this study is a correspondingly strong variation among cycad species in leaf physiology and structure. The values for leaf traits, and the coordination of traits showed considerable contrasts with other plant lineages. Most impressively, the cycads showed similar nutrient values, but lower gas exchange for given nutrient values, and no apparent coordination of hydraulics and gas exchange on a



Research 7

Fig. 4 Leaf photosynthetic nitrogen and phosphorus use efficiency (A_N and A_P , respectively), nutrient concentrations per mass (N_m , P_m , Fe_m , S_m , Zn_m , Cu_m) and chlorophyll concentration per mass (Chl_m) in relation to leaf mass per area (LMA) across cycad species in Xishuangbanna Tropical Botanical Garden (XTBG, green circles) and Fairylake Botanical Garden (FBG, red circles). Points are means \pm SE. Linear (a–f) or exponential regressions (g) were fitted to the data with shade-tolerant species (open circles) not included.

leaf area basis across species. However, the cycads were convergent with other lineages in their leaf economic spectrum relationships. A recent molecular dating study suggested that the extant cycads are not older than *c*. 12 Myr (Nagalingum *et al.*, 2011). However, we hypothesize that these trait relationships first evolved in the ancient cycads and remain in modern cycads because first, cycads show relative morphological, and potentially, physiological stability in the face of long-term climate change (Haworth *et al.*, 2011), and second, fossil and extant cycads have strong morphological similarities.

Just as studies have shown these relationships to hold within and across extant biomes (Reich *et al.*, 1997; Wright *et al.*, 2004), our study extends the existence of these relationships to a lineage dating back 250 Myr, suggesting that they potentially existed for



Fig. 5 The relationship between light-saturated photosynthetic rate and maximum leaf hydraulic conductance expressed on an area basis (A_a and K_{leaf-a} ; a), and on a mass basis (A_m and K_{leaf-m} ; b) and box plots of A_a (c) and K_{leaf-a} ; d) of cycads compared with other plant groups. Green circles, Xishuangbanna Tropical Botanical Garden (XTBG) cycads; red circles, Fairylake Botanical Garden (FBG) cycads; Blue open circles, shade-tolerant species. (a, b) Error bars indicate \pm SE. The solid line is a linear regression fitted to the data with shade-tolerant species (blue open circles) not included. The grey points are data for different taxonomic plant groups from Brodribb *et al.* (2007). (c, d) Boundaries of the boxes indicate the 25th and 75th percentiles, whiskers denote the 90th and 10th percentiles, points represent observations beyond these percentiles and internal lines denote the medians.



Fig. 6 Relationships between leaf mass per area (LMA) and mass based light-saturated photosynthetic rate (A_m), between LMA and leaf nitrogen concentration (N_m), and between N_m and A_m across cycads in Xishuangbanna Tropical Botanical Garden (XTBG, green circles) and Fairylake Botanical Garden (FBG, red circles) and the global dataset (Glopnet). The grey points are global dataset of Wright *et al.* (2004). All variables were log-transformed. Solid lines are standardized major axis (SMA) lines fitted to the global dataset, whereas break lines SMA lines fitted to the cycad species with shade-tolerant species (blue open circles) not included. ns, no significant difference (P > 0.05); *, P < 0.05; **, P < 0.01; ***, P < 0.001.

the species that dominated in those now extinct ecosystems. Further, the presence of these relationships among cycads just as previously found for angiosperms and conifers indicates that these trait relationships appeared multiple times as leaves evolved distinctly in different major seed plant lineages.

High variation in leaf functional traits and comparisons with other lineages

The high variation in leaf structural and physiological traits found among the cycad species within common gardens points to genetic divergences across species native to contrasting



Fig. 7 Relationships between maximum leaf hydraulic conductance expressed on an area basis (K_{leaf-a} ; a) and leaf mass per area (LMA), and between K_{leaf} on a mass basis (K_{leaf-m} ; b) and LMA across cycad species in Xishuangbanna Tropical Botanical Garden (XTBG, green circles) and Fairylake Botanical Garden (FBG, red circles). The grey points are a combined dataset mainly for angiosperms from Simonin *et al.* (2012). The solid line is a standardized major axis (SMA) line fitted to the combined angiosperm dataset, whereas the break line a SMA line fitted to the cycad species with shade-tolerant species (blue open circles) not included. ns, no significant difference (P > 0.05); **, P < 0.01.

environments (Monson, 1996). Although the dominance of cycads in terrestrial ecosystems has declined since the late Mesozoic, cycads still grow successfully in a wide range of habitats from very moist tropical rainforest to dry lands and high elevations (Norstog & Nicholls, 1997; Whitelock, 2002). The native habitats of the cycad species used in this study also varied greatly in their native water regimes. The cycads varied especially strongly in traits known to shift across habitats differing in water availability, such as LMA, photosynthetic traits and leaf hydraulic conductance (K_{leaf}) . Indeed, the range of values for cycads extended across the whole range of the values previously reported for other gymnosperms. Notably, cycads possessed higher $K_{\text{leaf-a}}$ and A_{a} than more primitive plant groups including bryophytes, lycophytes and ferns (Brodribb et al., 2007; Waite & Sack, 2010), which contributes to higher relative growth rates for a given level of leaf allocation (Sack et al., 2005; Zhang & Cao, 2009). Recently the higher maximum values of $K_{\text{leaf-a}}$ and A_{a} of angiosperms has been proposed to have contributed to, or driven their dominance over earlier groups since the Late Cretaceous (Brodribb & Feild, 2010; Feild et al., 2011a). Similarly, the higher values for cycads than modern members of more primitive lineages (bryophytes, mosses and ferns), together with their high diversification in leaf structure and function, provide a potential physiological explanation for the dominance of cycads across a variety of terrestrial habitats in the Mesozoic, assuming that this range of trait values also characterized ancestral cycads, as discussed earlier.



Fig. 8 Principal component analyses (PCA) of mean values of leaf traits for 33 cycad species. Trait (a) and species (b, c) loadings on the first and second axes. Trait symbols are in Supporting Information Table S2 and species codes in Table S3. (b) Green circles, cycads in Xishuangbanna Tropical Botanical Garden (XTBG); red circles, cycads in Fairylake Botanical Garden (FBG). (c) White circles, species from dry habitats; black circles, species from moist habitats; grey circles, space.

Compared with angiosperms, however, cycads, like other gymnosperms, exhibited lower maximum and/or mean values for $K_{\text{leaf-a}}$, A_{a} and A_{m} (Table 1). These traits would be related to low relative growth rates and lead to a major disadvantage for gymnosperms compared with angiosperms (Bond, 1989; Berendse & Scheffer, 2009). The lower $K_{\text{leaf-a}}$ of the cycads compared with angiosperms may relate to their low leaf vein density, which limits development of high transpiration and photosynthetic rates and imposed a major disadvantage on cycads competing with angiosperms (Boyce *et al.*, 2009). Indeed, cycads had lower A_m for a given N_m than the mean for seed plants (Fig. 6c), that is, lower photosynthetic N use efficiencies (A_N) . This difference may have arisen due to greater partitioning of N to nonphotosynthetic structures (Hikosaka, 2004). Although information on N partitioning of cycads is unavailable, cycads have high concentrations of lignin, leaf mucilage and other secondary compounds (Norstog & Nicholls, 1997; Brenner et al., 2003), requiring substantial

investment of N and other nutrients. The higher LMA of cycads compared with the Glopnet mean agrees with their high investments in leaf toughness and herbivory defense (Clark & Clark, 1988; Prado *et al.*, 2014) and potentially high leaf longevity compared with other seed plant lineages (up to 106 months for *M. riedlei*; Wright *et al.*, 2004). The lower $K_{\text{leaf-a}}$, A_{a} , A_{m} and A_{N} of cycads than angiosperms would impose disadvantages for competing with later seed plants, and by hypothesis, would have contributed to the decline in dominance of cycads in late Mesozoic (Cretaceous), when angiosperms began to diversify and establish dominance (Crane *et al.*, 1995; Berendse & Scheffer, 2009).

Relationship between leaf hydraulic conductance and leaf photosynthetic capacity in cycads

A positive relationship between area-based $K_{\text{leaf-a}}$ and A_{a} previously has been shown for plants across a variety of taxonomic groups including bryophytes, lycopods, ferns, gymnosperms and angiosperms (Brodribb et al., 2007). As described in the Introduction, the close relationship between leaf hydraulics and photosynthesis has been generally explained by the importance of efficient leaf water supply for maintaining relatively high leaf water potential during transpiration to allow stomata to remain open. However, no association was found between $K_{\text{leaf-a}}$ and $A_{\rm a}$ across the cycad species studied. The absence of a relationship between $K_{\text{leaf-a}}$ and A_{a} may be due to the fact that the cycad species originate in habitats with a wide range of water availabilities (e.g. soil water potential and vapor pressure deficit). Plants of a given life form and from habitats with similar water availabilities converge in a relatively narrow range of $\Psi_{\rm L}$ during peak transpiration, and therefore gs could be associated with K (hydraulic conductance) as $g_s = K \times (\Psi_L - \Psi_{soil})/VPD$. Species from habitats with different water availabilities may operate at different $\Psi_{\rm L}$, decoupling the coordination between leaf hydraulic conductance and gas exchange properties (Sack et al., 2005; Feild et al., 2011b; Sack & Scoffoni, 2013). However, no coordination between leaf water transport and photosynthetic rate was detected even when the relationship was tested in species native to habitats with similar water availabilities (data not shown). Alternatively, decoupling between leaf water transport and photosynthetic rate could have arisen due to the influence of other factors such as leaf water storage, especially as cycad leaves possess hypodermis and transfusion tissues (Hu & Yao, 1981), which might allow stomata to remain open despite low water transport across the lamina. Indeed, cycads tend to have higher A_a despite similar $K_{\text{leaf-a}}$ to conifers (Table 1; Fig. 5), representing a higher photosynthetic performance relative to hydraulic investment. Another potential factor contributing to the water and carbon decoupling is that some cycad species could be Crassulacean acid metabolism (CAM)-facultative plants (e.g. D. edule) (Vovides et al., 2002), which would result in relatively high photosynthetic water use efficiencies. The relationship between $K_{\text{leaf-m}}$ and A_{m} would arise from a negative relationship between K_{leaf-m} and LMA (Fig. 7b). Indeed, the negative relationship between $K_{\text{leaf-m}}$ and LMA in cycads generally agreed with the relationship reported

for angiosperms (Simonin *et al.*, 2012), suggesting a strong generality in this relationship. The mathematical source of this negative relationship is that $K_{\text{leaf-m}}$ is calculated as $K_{\text{leaf-a}}$ divided by LMA, and that $K_{\text{leaf-a}}$ is independent of LMA. The significance of the trend is a trade-off between mass investment in leaf hydraulics and in leaf longevity (Simonin *et al.*, 2012). The lower $K_{\text{leaf-m}}$ at a given LMA in cycads than for angiosperms suggests a relatively low resource investment in leaf hydraulic system.

Trait relationships in cycads with respect to the global leaf economic spectrum

The relationships of fundamental leaf functional traits for cycads generally agreed with previously reported trait relationships in the global leaf economic spectrum (Field & Mooney, 1986; Reich et al., 1997; Wright et al., 2004). Recent studies have also revealed that ferns and mosses follow the global leaf economic spectrum in general, though LMA was replaced by canopy mass per area for mosses (Karst & Lechowicz, 2007; Waite & Sack, 2010). Therefore, the fundamental ecological and physiological trade-offs constraining leaf functional traits apply not only to late-derived seed plants, but also to relatively primitive living plants such as mosses, ferns and cycads, confirming the generality of the principal trade-offs. Notably, Z. fischeri showed much lower LMA and leaf thickness, and much higher Am, Kleaf-m and nutrient concentrations than other cycad species. However, the leaf traits of even this 'outlier' species followed most of the functional trait relationships among other species, indicating tightly bounded domains constraining leaf traits (Reich et al., 1997; Meinzer, 2003; Shipley et al., 2006).

Another novel aspect of our study is the expanded set of nutrient relationships important in the leaf economic spectrum. In addition to the relationships among LMA, A_m, N_m and P_m, we found relationships of LMA and A_m with Fe_m and S_m, as well as between LMA and photosynthetic nutrient use efficiencies (A_N and $A_{\rm P}$). Indeed, Fe_m was more strongly related to $A_{\rm m}$ than N_m, which agrees with the Chinese traditional knowledge that cycads show strong preference for Fe and soil addition of Fe may increase growth rates of some cycads. Fe is related to numerous electron transfer reactions and biochemical processes in photosynthesis and is required for biosynthesis of chlorophyll (e.g. Spiller & Terry, 1980). The concentration of Fe in the nontoxic range has been positively related to the amount of photosynthetic machinery and consequently to photosynthetic capacity (Spiller & Terry, 1980). The generality of these relationships needs to be tested in other taxonomic groups. Notably, the nutrient concentrations of cycads were close to the mean values for dominant angiosperm species across geographical locations in China (Hou, 1982), whereas N_m and P_m of cycads were even slightly higher than the Glopnet mean. Although cycads are often able to fix N via symbiotic cyanobacteria in roots, the strong relationship between nutrient concentrations and Am suggests a conserved proportion of nutrients invested in photosynthetic machinery and thus to $A_{\rm m}$. A limitation imposed by the amount of photosynthetic machinery on $A_{\rm m}$ was also indicated by the strong

relationship between Chl_m and A_m . Apparently due to a lack of data of nutrient concentrations other than N and P in the global dataset, interspecific relationships among concentrations of nutrients other than N and P with A_m and LMA have not been established across other taxonomic groups.

In conclusion, the relationships among cycad leaf functional traits generally agreed with the global economic spectrum, confirming the generality of the fundamental trade-offs, and potentially extending these principles of leaf design to the ancient past. High leaf hydraulic conductance and photosynthetic rate in cycads compared with bryophytes, lycophytes and ferns, as well as high diversification in leaf structure and function would have facilitated their dominance in a variety of ecosystems in the Mesozoic. Low leaf water supply and photosynthetic nutrient use efficiencies in cycads relative to angiosperms could be a potential factor in limiting their development of higher growth rates. Nutrients other than N (e.g. Fe and S) explained a significant amount of variation in Am of cycads as well, suggesting the importance of introducing nutrients other than N and P into a higher resolution understanding of the leaf economic spectrum. Interestingly, no coordination between leaf hydraulic conductance and photosynthesis was found among cycads but overall they tended to have higher photosynthetic performance relative to hydraulic capacity compared with other plant lineages.

Acknowledgements

We thank Dr D. Ackerly and three anonymous reviewers for their careful reading of our manuscript, and their constructive comments and suggestions. We thank the Biogeochemistry Laboratory of our botanical garden for the determination of nutrient concentrations. We also would like to thank Mrs Fu Xuewei, Yang Qiuyun, Zeng Xiaodong and Li Shuai for their assistance in the field measurements, and S. Mathews and M. Bartlett for assistance with analyses. Support from the National Natural Science Foundation of China (31170399) is gratefully acknowledged.

References

- Ares A, Burner DM, Brauer DK. 2009. Soil phosphorus and water effects on growth, nutrient and carbohydrate concentrations, delta C¹³, and nodulation of mimosa (*Albizia julibrissin* Durz.) on a highly weathered soil. *Agroforestry Systems* 76: 317–325.
- Berendse F, Scheffer M. 2009. The angiosperm radiation revisited, an ecological explanation for Darwin's 'abominable mystery'. *Ecology Letters* 12: 865–872.
- Blonder B, Violle C, Bentley LP, Enquist BJ. 2011. Venation networks and the origin of the leaf economics spectrum. *Ecology Letters* 14: 91–100.
- Blonder B, Violle C, Bentley LP, Enquist BJ. 2014. Inclusion of vein traits improves predictive power for the leaf economic spectrum: a response to Sack *et al.* (2013). *Journal of Experimental Botany* 65: 5109–5114.
- Blonder B, Violle C, Enquist BJ. 2013. Assessing the causes and scales of the leaf economics spectrum using venation networks in Populus tremuloides. *Journal* of Ecology 101: 981–989.
- Bond WJ. 1989. The tortoise and the hare: ecology of angiosperm dominance and gymnosperm persistence. *Biological Journal of the Linnean Society* 36: 227– 249.
- Boyce CK, Brodribb TJ, Feild TS, Zwieniecki MA. 2009. Angiosperm leaf vein evolution was physiologically and environmentally transformative. *Proceedings* of the Royal Society B 276: 1771–1776.

- Brenner ED, Stevenson DW, Twigg RW. 2003. Cycads: evolutionary innovations and the role of plant-derived neurotoxins. *Trends in Plant Science* 8: 446–452.
- Brodribb TJ, Feild TS. 2010. Leaf hydraulic evolution led a surge in leaf photosynthetic capacity during early angiosperm diversification. *Ecology Letters* 13: 175–183.
- Brodribb TJ, Feild TS, Jordan GJ. 2007. Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Physiology* 144: 1890–1898.
- Campanello PI, Gatti MG, Goldstein G. 2008. Coordination between water-transport efficiency and photosynthetic capacity in canopy tree species at different growth irradiances. *Tree Physiology* 28: 85–94.
- Castelli F, Contillo R, Miceli F. 1996. Non-destructive determination of leaf chlorophyll content in four crop species. *Journal of Agronomy and Crop Science* 177: 275–283.
- Chaw SM, Walters TW, Chang CC, Hu SH, Chen SH. 2005. A phylogeny of cycads (Cycadales) inferred from chloroplast *matK* gene, *trnK* intron, and
- nuclear rDNA ITS region. *Molecular Phylogenetics and Evolution* 37: 214–234. Clark DB, Clark DA. 1988. Leaf production and the cost of reproduction in the
- neotropical rainforest cycad, *Zamia skinneri. Journal of Ecology* 76: 1153–1163. Cramer MD, Hawkins HJ, Verboom GA. 2009. The importance of nutritional regulation of plant water flux. *Oecologia* 161: 15–24.
- Crane PR, Friis EM, Pedersen KR. 1995. The origin and early diversification of angiosperms. *Nature* 374: 27–33.
- Fanizza G, Gatta CD, Bagnulo C. 1991. A non-destructive determination of leaf chlorophyll in Vitis vinifera. The Annals of Applied Biology 119: 203–205.
- Feild TS, Brodribb TJ, Iglesias A, Chatelet DS, Baresch A, Upchurch GR, Gomez B, Mohr BAR, Coiffard C, Kvacek J et al. 2011a. Fossil evidence for Cretaceous escalation in angiosperm leaf vein evolution. Proceedings of the National Academy of Sciences, USA 108: 8363–8366.
- Feild TS, Upchurch GR, Chatelet DS, Brodribb TJ, Grubbs KC, Samain MS, Wanke S. 2011b. Fossil evidence for low gas exchange capacities for Early Cretaceous angiosperm leaves. *Paleobiology* 37: 195–213.
- Field C, Mooney HA. 1986. The photosynthesis–nitrogen relationship in wild plants. In: Givnish T, ed. On the economy of plant form and function. London, UK: Cambridge University Press, 25–55.
- Gao Z, Thomas BA. 1989. A review of fossil cycad megasporophylls, with new evidence of crossozamia pomel and its associated leaves from the Lower Permian of Taiyuan, China. *Review of Palaeobotany and Palynology* 60: 205–223.
- Gratani L. 1992. A nondestructive method to determine chlorophyll content of leaves. *Photosynthetica* 26: 469–473.
- Haworth M, Fitzgerald A, McElwain JC. 2011. Cycads show no stomatal-density and index response to elevated carbon dioxide and subambient oxygen. *Australian Journal of Botany* 59: 629–638.
- Hikosaka K. 2004. Interspecific difference in the photosynthesis–nitrogen relationship: patterns, physiological causes, and ecological importance. *Journal of Plant Research* 117: 481–494.
- Hou XY. 1982. Chinese vegetable geography and chemical elements: analyses of the dominant plant species. Beijing, China: Science Press.
- Hu Y-S, Yao B-J. 1981. Transfusion tissue in gymnosperm leaves. *Botanical Journal of the Linnean Society* 83: 263–272.
- Huang Y-Y, Zhang H-D. 1999. The brief report on first discovery of vessel in cycads. *Journal of Agricultural and Biological Science* 18: 99–100.
- Karst AL, Lechowicz MJ. 2007. Are correlations among foliar traits in ferns consistent with those in the seed plants? *New Phytologist* 173: 306–312.
- Kuppers M. 1984. Carbon relations and competition between woody species in a central European hedgerow. 2. Stomatal responses, water-use, and hydraulic conductivity in the root leaf pathway. *Oecologia* 64: 344–354.
- Lloyd J, Bloomfield K, Domingues TF, Farquhar GD. 2013. Photosynthetically relevant foliar traits correlating better on a mass vs an area basis: of ecophysiological relevance or just a case of mathematical imperatives and statistical quicksand? *New Phytologist* 199: 311–321.
- Mamay SH. 1969. Cycads: fossil evidence of late Paleozoic origin. *Science* 164: 295–296.
- Manetas Y, Grammatikopoulos G, Kyparissis A. 1998. The use of the portable, non-destructive, SPAD-502 (Minolta) chlorophyll meter with leaves of varying

trichome density and anthocyanin content. *Journal of Plant Physiology* **153**: 513–516.

Marenco RA, Antezana-Vera SA, Nascimento HCS. 2009. Relationship between specific leaf area, leaf thickness, leaf water content and SPAD-502 readings in six Amazonian tree species. *Photosynthetica* 47: 184–190.

Markwell J, Osterman JC, Mitchell JL. 1995. Calibration of the Minolta SPAD-502 leaf chlorophyll meter. *Photosynthesis Research* 46: 467–472.

Marquard R, Tipton J. 1987. Relationship between extractable chlorophyll and an in situ method to estimate leaf greenness. *HortScience* 22: 1327.

Meinzer FC. 2003. Functional convergence in plant responses to the environment. *Oecologia* 134: 1–11.

Meinzer FC, Grantz DA. 1990. Stomatal and hydraulic conductance in growing sugarcane – stomatal adjustment to water transport capacity. *Plant, Cell & Environment* 13: 383–388.

Melcher PJ, Michele Holbrook N, Burns MJ, Zwieniecki MA, Cobb AR, Brodribb TJ, Choat B, Sack L. 2012. Measurements of stem xylem hydraulic conductivity in the laboratory and field. *Methods in Ecology and Evolution* 3: 685–694.

Mencuccini M. 2003. The ecological significance of long-distance water transport: short-term regulation, long-term acclimation and the hydraulic costs of stature across plant life forms. *Plant, Cell & Environment* 26: 163–182.

Monson RK. 1996. The use of phylogenetic perspective in comparative plant physiology and development biology. *Annals of the Missouri Botanical Garden* 83: 3–16.

Nagalingum NS, Marshall CR, Quental TB, Rai HS, Little DP, Mathews S. 2011. Recent synchronous radiation of a living fossil. *Science* 334: 796–799.

Niinemets Ü, Sack L. 2006. Structural determinants of leaf light-harvesting capacity and photosynthetic potentials. In: Esser K, Lüttge U, Beyschlag W, Murata J, eds. *Progress in botany.* Berlin/Heidelberg, Germany: Springer, 385–419.

Norstog K, Nicholls TJ. 1997. *The biology of the cycads*. Ithaca, NY, USA: Comstock Publishing Associates/ Cornell University Press.

Osnas JLD, Lichstein JW, Reich PB, Pacala SW. 2013. Global leaf trait relationships: mass, area, and the leaf economics spectrum. *Science* 340: 741–744.

Parkhurst DF. 1994. Diffusion of CO₂ and other gases inside leaves. *New Phytologist* **126**: 449–479.

Pearson K. 1901. On lines and planes of closest fit to systems of points in space. *Philosophical Magazine* 2: 559–572.

Prado A, Sierra A, Windsor D, Bede JC. 2014. Leaf traits and herbivory levels in a tropical gymnosperm, *Zamia stevensonii* (Zamiaceae). *American Journal of Botany* 101: 437–447.

Randall PJ, Bouma D. 1973. Zinc deficiency, carbonic-anhydrase, and photosynthesis in leaves of spinach. *Plant Physiology* 52: 229–232.

Reich PB, Ellsworth DS, Walters MB. 1998. Leaf structure (specific leaf area) modulates photosynthesis–nitrogen relations: evidence from within and across species and functional groups. *Functional Ecology* **12**: 948–958.

Reich PB, Walters MB, Ellsworth DS. 1997. From tropics to tundra: global convergence in plant functioning. *Proceedings of the National Academy of Sciences, USA* 94: 13730–13734.

Roderick ML, Berry SL, Noble IR, Farquhar GD. 1999a. A theoretical approach to linking the composition and morphology with the function of leaves. *Functional Ecology* 13: 683–695.

Roderick ML, Berry SL, Saunders AR, Noble IR. 1999b. On the relationship between the composition, morphology and function of leaves. *Functional Ecology* 13: 696–710.

Sack L, Frole K. 2006. Leaf structural diversity is related to hydraulic capacity in tropical rain forest trees. *Ecology* 87: 483–491.

Sack L, Melcher PJ, Zwieniecki MA, Holbrook NM. 2002. The hydraulic conductance of the angiosperm leaf lamina: a comparison of three measurement methods. *Journal of Experimental Botany* 53: 2177–2184.

Sack L, Scoffoni C. 2012. Measurement of leaf hydraulic conductance and stomatal conductance and their responses to irradiance and dehydration using the evaporative flux method (EFM). *Journal of Visualized Experiments: JoVE* 70: e4179.

Sack L, Scoffoni C, John GP, Poorter H, Mason CM, Mendez-Alonzo R, Donovan LA. 2013. How do leaf veins influence the worldwide leaf economic spectrum? Review and synthesis. *Journal of Experimental Botany* 64: 4053– 4080.

Sack L, Scoffoni C, John GP, Poorter H, Mason CM, Mendez-Alonzo R, Donovan LA. 2014. Leaf mass per area is independent of vein length per area: avoiding pitfalls when modelling phenotypic integration (reply to Blonder *et al.* 2014). *Journal of Experimental Botany* 65: 5115–5123.

Sack L, Tyree MT, Holbrook NM. 2005. Leaf hydraulic architecture correlates with regeneration irradiance in tropical rainforest trees. *New Phytologist* 167: 403–413.

Sage RF, Pearcy RW, Seemann JR. 1987. The nitrogen use efficiency of C_3 and C_4 plants III. Leaf nitrogen effects on the activity of carboxylating enzymes in *Chenopodium album* (L.) and *Amaranthus retroflexus* (L.). *Plant Physiology* 85: 355–359.

Santiago LS, Goldstein G, Meinzer FC, Fisher JB, Machado K, Woodruff D, Jones T. 2004. Leaf photosynthetic traits scale with hydraulic conductivity and wood density in Panamanian forest canopy trees. *Oecologia* 140: 543–550.

Shipley B, Lechowicz MJ, Wright I, Reich PB. 2006. Fundamental trade-offs generating the worldwide leaf economics spectrum. *Ecology* 87: 535–541.

Simonin KA, Limm EB, Dawson TE. 2012. Hydraulic conductance of leaves correlates with leaf lifespan: implications for lifetime carbon gain. *New Phytologist* 193: 939–947.

Singha A, Townsend E. 1989. Relationship between chromaticity values and chlorophyll concentration in apple, grape, and peach leaves. *HortScience* 24: 1034.

Sober A. 1997. Hydraulic conductance, stomatal conductance, and maximal photosynthetic rate in bean leaves. *Photosynthetica* **34**: 599–603.

Spiller S, Terry N. 1980. Limiting factors in photosynthesis: II. iron stress diminishes photochemical capacity by reducing the number of photosynthetic units. *Plant Physiology* 65: 121–125.

Stevenson DW, Norstog KJ, Molsen DV. 1996. Midribs of cycad pinnae. *Brittonia* 48: 67–74.

Uddling J, Gelang-Alfredsson J, Piikki K, Pleijel H. 2007. Evaluating the relationship between leaf chlorophyll concentration and SPAD-502 chlorophyll meter readings. *Photosynthesis Research* **91**: 37–46.

Vovides AP, Etherington JR, Dresser PQ, Groenhof A, Iglesias C, Ramirez JF. 2002. CAM-cycling in the cycad *Dioon edule* Lindl. in its natural tropical deciduous forest habitat in central Veracruz, Mexico. *Botanical Journal of the Linnean Society* 138: 155–162.

Waite M, Sack L. 2010. How does moss photosynthesis relate to leaf and canopy structure? Trait relationships for 10 Hawaiian species of contrasting light habitats. *New Phytologist* 185: 156–172.

Warton DI, Wright IJ, Falster DS, Westoby M. 2006. Bivariate line-fitting methods for allometry. *Biological Reviews of the Cambridge Philosophical Society* 81: 259–291.

Westoby M, Reich PB, Wright IJ. 2013. Understanding ecological variation across species: area-based vs mass-based expression of leaf traits. *New Phytologist* 199: 322–323.

Whitelock LM. 2002. The Cycads. Portland, OR, USA: Timber Press.

Witkowski ETF, Lamont BB. 1991. Leaf specific mass confounds leaf density and thickness. *Oecologia* 88: 486–493.

Wright IJ, Reich PB, Westoby M. 2003. Least-cost input mixtures of water and nitrogen for photosynthesis. *The American Naturalist* 161: 98–111.

Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F, Cavender-Bares J, Chapin T, Cornelissen JHC, Diemer M et al. 2004. The worldwide leaf economics spectrum. *Nature* 428: 821–827.

Zgurski JM, Rai HS, Fai QM, Bogler DJ, Francisco-Ortega J, Graham SW. 2008. How well do we understand the overall backbone of cycad phylogene? New insights from a large, multigene plastid data set. *Molecular Phylogenetics and Evolution* 47: 1232–1237.

Zhang JL, Cao KF. 2009. Stem hydraulics mediates leaf water status, carbon gain, nutrient use efficiencies and plant growth rates across dipterocarp species. *Functional Ecology* 23: 658–667.

Supporting Information

Additional supporting information may be found in the online version of this article.

Table S1 Cycad species studied in Xishuangbanna TropicalBotanical Garden (XTBG) and Fairylake Botanical Garden(FBG), origin and native habitats

Table S2 Traits, symbols and units

Table S3 Means and SE for each trait by species

Please note: Wiley Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



• For submission instructions, subscription and all the latest information visit www.newphytologist.com